

# Prevalence of depression among female adolescents in Indonesia: A cross-sectional study using data from the 2023 Indonesian Health Survey (SKI)

Zahra N. Syifa<sup>1\*</sup>, Vernonia Yora Saki<sup>2\*</sup>, Ahmad Azmi Nasution<sup>2\*</sup>, Rizkianti Anggraini<sup>3</sup>, Aulia Rahman<sup>4</sup>

## AFFILIATION

**1** Undergraduate Medical Program, Faculty of Medicine and Health Sciences, Universitas Bengkulu, Bengkulu, Indonesia

**2** Department of Community Medicine, Faculty of Medicine and Health Sciences, Universitas Bengkulu, Bengkulu, Indonesia

**3** Department of Anatomy, Faculty of Medicine and Health Sciences, Universitas Bengkulu, Bengkulu, Indonesia

**4** Department of Public Health, Faculty of Public Health, Universitas Andalas, Padang, Indonesia

\*Contributed equally

## CORRESPONDENCE TO

Vernonia Yora Saki. Department of Community Medicine, Faculty of

Popul. Med. 2026;8(2):8

Medicine and Health Sciences, Universitas Bengkulu, Jl. WR. Supratman, Kandang Limun, Kec. Muara Bangka Hulu, Sumatera, 38371, Bengkulu, Indonesia

E-mail: [yorasaki@unib.ac.id](mailto:yorasaki@unib.ac.id)

ORCID iD: <https://orcid.org/0000-0003-0606-6919>

## KEYWORDS

adolescent, depression, marital status, smoking status, urban residence

**Received:** 16 February 2026, **Revised:** 10 April 2026,

**Accepted:** 21 April 2026

<https://doi.org/10.18332/popmed/220962>

## ABSTRACT

**INTRODUCTION** In Indonesia, evidence specifically focused on depression among female adolescents aged 15–19 years is still limited. This study aimed to determine the prevalence of depression and its associated factors among adolescent girls aged 15–19 years in Indonesia using data from the 2023 Indonesian Health Survey (SKI).

**METHODS** This cross-sectional study used secondary data from SKI 2023. The analytic sample included 26554 female adolescents aged 15–19 years. Depression was assessed using the Mini International Neuropsychiatric Interview (MINI). Smoking status was categorized as never smoker, ex-smoker, and current smoker. Covariates included employment status, marital status, education level, and location of residence. Data were analyzed using chi-squared tests and multiple logistic regression accounting for the complex survey design.

**RESULTS** The prevalence of depression was 1.6%. Smoking status was significantly associated with depression in bivariate analysis ( $p=0.001$ ). In multivariable analysis, ex-smokers had higher odds of depression than never smokers (AOR=10.93; 95% CI: 10.73–11.13), followed by current smokers (AOR=5.82; 95% CI: 5.58–6.08). Divorced/widowed adolescents had higher odds of depression than those who had never married (AOR=4.32; 95% CI: 4.15–4.49). Adolescents living in an urban location also had higher odds of depression than those living in a rural location (AOR=2.12; 95% CI: 2.10–2.13).

**CONCLUSIONS** Depression affected 1.6% of female adolescents aged 15–19 years in Indonesia. Smoking status, marital status, and location of residence were associated with depression in this population.

## INTRODUCTION

Mental health is an important aspect of adolescent development. This period is a transitional phase characterized by very rapid biological, psychological, and social changes that make it susceptible to the emergence of various mental health problems<sup>1-3</sup>. The World Health Organization (WHO) confirms that around 1 in 7 adolescents in the world experience mental disorders, including depression<sup>2</sup>. Depression in adolescence is one of the most impactful health problems, as it is associated with poorer academic and social functioning in the future, which suggests

that the impact of depression on adolescence can extend far beyond the adolescent period itself<sup>4,5</sup>.

The burden of mental health problems in adolescents in Indonesia also shows a tendency that needs attention. I-NAMHS reports that about 1 in 3 adolescents (34.9%) have mental health problems, and about 1 in 20 adolescents (5.5%) have met the criteria for a mental disorder<sup>6</sup>. In addition, I-NAMHS also showed that the prevalence of depression in female adolescents (6.7%) was higher than that of adolescent boys (4.0%), which indicates a gender-based vulnerability to adolescent mental health problems<sup>6</sup>.

In line with that, the 2023 Indonesian Health Survey (SKI) reported that the national prevalence of depression was 1.4%, with the highest proportion of 2.0% found in the age group of 15–24 years<sup>7</sup>. In this young age group, the prevalence of depression in women (2.8%) is also higher than in men (1.1%), which further confirms that female adolescents are a more vulnerable group to depression<sup>7</sup>.

Previous research related to depression in Indonesia has found that sociodemographic factors, including gender, marital status, location of residence, and behavioral factors such as smoking, contribute to depression in adolescents<sup>8,9</sup>. Previous research has shown that female adolescents have a higher risk of depression than adolescent boys<sup>8</sup>. In line with these findings, this study aims to determine the prevalence of depression and factors associated with it in female adolescents aged 15–19 years in Indonesia using the 2023 SKI Survey.

## METHODS

### Study design and data source

We conducted a secondary dataset analysis of the cross-sectional Indonesian Health Survey 2023. This data collection was conducted nationwide by the National Development Planning Agency (BKPK) from August to October 2023, covering 38 provinces and 514 districts/cities in Indonesia. Therefore, these data can represent the health status at the national to district/city level<sup>7</sup>.

### Participants

The Indonesian Health Survey covered the entire territory of Indonesia, recording 1082958 household members, of whom with 877531 individuals completed the interview and measurement process<sup>7</sup>. In this study, participants for analysis were selected from the 2023 SKI individual data set using the inclusion criterion of being aged 15–19 years. After selection, 53845 respondents met these criteria. A total of 27291 male

respondents were excluded from the analysis, leaving a final analytical sample that consisted of 26554 female adolescents aged 15–19 years in Indonesia (Figure 1).

### Data collection


Data collection in the 2023 Indonesian Health Survey was conducted through interviews, measurements, and examinations. For regular households, the interview component used two instruments, namely the household questionnaire and the individual questionnaire. The dependent and independent variables analyzed in this study were obtained from the individual questionnaire in Block XI, which consists of Blocks A to J. The independent variable, smoking status, was derived from Block G, which covers knowledge and behavior. The dependent variable, depression, was obtained from Block C, which addresses mental health. Finally, covariates such as employment status, marital status, education level, and location of residence were obtained from the household questionnaire, specifically from Block IV (Household Member Information) and Block I (Location Identification)<sup>7</sup> (Supplementary file).

### Variables

#### Dependent variable

This study defines depression as a dependent variable. The depression variable was assessed using the Mini International Neuropsychiatric Interview (MINI) instrument, referring to symptoms experienced during the previous two weeks. The instrument consists of 10 questions with 'yes' or 'no' responses. Examples of the items include questions about persistent sadness over the previous two weeks, diminished interest or enjoyment in usual activities, and ongoing feelings of tiredness or low energy. Respondents were considered to have depression if they answered 'yes' to at least 2 of the 3 main questions (questions 1–3) and also 'yes' to at least 2 of the 7 additional questions (questions 4–10)<sup>10</sup>.

**Figure 1. Flowchart of participant selection for female adolescents aged 15-19 years included in the study, based on the Indonesian Health Survey 2023 (N=26554)**



*Independent variable*

Smoking status was defined as an independent variable in this study and classified into three categories: never smoker, ex-smoker, and current smoker. Respondents were categorized as never smokers if they had never tried smoking until the time of data collection. The ex-smoker category referred to respondents who had smoked previously, either daily or occasionally, but not on a daily basis at the time of assessment. Meanwhile, respondents were classified as current smokers if they had smoked every day or occasionally during the past month<sup>7</sup>.

**Covariates**

Covariates in this study included employment status, marital status, education level, and location of residence. Employment status was categorized as either employed or not employed. Marital status was classified as never married, married, or divorced/widowed. Education level was determined based on the highest level of formal education completed and grouped into at least high school education and lower than high school education. Location of residence was defined according to the village/urban ward classification and categorized as either urban or rural.

Covariates were selected based on prior literature indicating that these variables may be associated with depression<sup>8,9,11,12</sup>. Therefore, they were considered important for adjustment in the multivariable analysis using a determinant modeling approach, in which all variables were assumed to have an equal opportunity to influence the occurrence of depression. The selection of these covariates was based on Lawrence Green’s theory.

**Data management and statistical analysis**

The data management process began with extracting relevant variables from the 2023 Indonesian Health Survey database. Data cleaning was carried out to identify missing values and ensure logical consistency between sociodemographic variables and mental health instrument scores. The dependent variable, namely depression, was dichotomized using the Mini International Neuropsychiatric Interview (MINI) instrument at the cutoff point. Independent variables such as smoking status, age group, education level, employment status, marital status, and location of residence (urban, rural) were recoded to facilitate interpretation. The entire data processing process integrated weighting variables, strata, and primary sampling units (PSUs) to compensate for the complex survey design, enabling the estimation results to be generalized to the national population of female adolescents in Indonesia.

Data analysis was conducted in stages using an inferential statistical approach. Univariate analysis was used to describe the frequency distributions of respondents’ characteristics, with percentages reported for categorical variables and means ± standard deviations (SD) for continuous variables. Bivariate analysis was performed to test the relationship

between the independent variables and depression using the chi-squared test, accounting for the survey design. Furthermore, multivariate analysis was performed using multiple logistic regression to identify the most influential determinants of depression in female adolescents, after controlling for covariates. The results of the multivariate analysis were presented as crude odds ratios (OR) and adjusted odds ratios (AOR) with 95% confidence intervals. All statistical procedures were performed using software that supports the complex-samples module to ensure accurate standard error estimates.

**Table 1. Sociodemographic characteristics, age of smoking initiation, smoking status, and depressive symptoms among female adolescents aged 15–19 years in Indonesia, based on SKI 2023 (N=26554)**

Characteristics	n (%)
<b>Age (years)</b>	
15	5762 (21.7)
16	5895 (22.2)
17	5444 (20.5)
18	4806 (18.1)
19	4647 (17.6)
<b>Age (years), mean ± SD</b>	17.14 ± 1.34
<b>Education level</b>	
At least high school education	5523 (20.8)
Lower than high school education	21031 (79.2)
<b>Employment status</b>	
Employed	19677 (74.1)
Not employed	6877 (25.9)
<b>Marital status</b>	
Never married	25465 (95.9)
Divorced/widowed	53 (0.2)
Married	1036 (3.9)
<b>Residence</b>	
Rural	11206 (42.2)
Urban	15348 (57.8)
<b>Age of smoking initiation (years), mean ± SD</b>	15.48 ± 2.08
<b>Smoking status</b>	
Never smoker	26395 (99.4)
Ex-smoker	106 (0.4)
Current smoker	53 (0.2)
<b>Depression</b>	
No	26129 (98.4)
Yes	425 (1.6)

**Ethical aspects**

This study is a further analysis using secondary data from the 2023 Indonesian Health Survey (SKI). The survey protocol has been declared ethically sound by the Formal Health Research Ethics Commission of the Health Development Policy Agency (BKPK), Indonesian Ministry of Health, under number LB.02.01/2/KE.024/2023. All respondent identities in the dataset have been anonymized by the survey organizers to maintain the confidentiality of the research subjects.

**RESULTS**

Table 1 presents the distribution of sociodemographic characteristics, age of smoking initiation, smoking status,

and depression among 26554 female adolescents aged 15–19 years in Indonesia. The largest age groups were aged 16 years (22.2%) and aged 15 years (21.7%), with a mean age of  $17.14 \pm 1.34$  years. Most participants had lower than a high school education (79.2%). The majority were employed (74.1%), while 25.9% were not employed. Most female adolescents had never been married (95.9%), followed by those who were married (3.9%) and divorced/widowed (0.2%). Based on the location of residence, more participants lived in urban areas (57.8%) than in rural areas (42.2%). The mean age of smoking initiation was  $15.48 \pm 2.08$  years. Most participants had never smoked (99.4%), while 0.4% reported being ex-smokers and 0.2% reported being current smokers. The prevalence of depression in this study was 1.6% among

**Table 2. Bivariate analysis of smoking status and depressive symptoms using chi-squared test among female adolescents aged 15–19 years, based on SKI 2023 (N=26554)**

Smoking status	Depression		Total n	p	OR	95% CI:
	No n (%)	Yes n (%)				
Total, n	25723	831	26554			
Never smoker (ref.)	25603 (97.0)	792 (3.0)	26395		1	
Ex-smoker	77 (72.6)	29 (27.4)	106	0.001	12.59	12.37–12.82
Current smoker	43 (81.1)	10 (18.9)	53	0.001	8.10	7.79–8.43

**Table 3. Adjusted prevalence odds ratios (AOR) for depressive symptoms by smoking status and sociodemographic covariates among female adolescents aged 15–19 years, SKI 2023 (N=26554)**

Variable	OR (95% CI)	AOR (95% CI)
<b>Smoking status</b>		
Never smoker (ref.)	1	1
Ex-smoker	12.59 (12.37–12.82)	10.93 (10.73–11.13)
Current smoker	8.10 (7.79–8.43)	5.82 (5.58–6.08)
<b>Education level</b>		
At least high school education (ref.)	1	1
Lower than high school education	0.92 (0.91–0.93)	1.04 (1.03–1.05)
<b>Residence</b>		
Rural (ref.)	1	1
Urban	2.15 (2.13–2.16)	2.12 (2.10–2.13)
<b>Employment status</b>		
Employed (ref.)	1	1
Not employed	1.22 (1.21–1.23)	1.27 (1.26–1.28)
<b>Marital status</b>		
Never married (ref.)	1	1
Divorced/widowed	4.98 (4.80–5.16)	4.32 (4.15–4.49)
Married	0.85 (0.84–0.87)	0.81 (0.80–0.83)

The results of the multivariate analysis were presented as crude odds ratios (OR) and adjusted odds ratios (AOR) with 95% confidence intervals.

female adolescents aged 15–19 years in Indonesia.

The prevalence of depression was higher among ex-smokers (27,4%), followed by current smokers (18,9%). Bivariate analysis with chi-squared test showed a significant association between smoking status and depression ( $p=0.001$ ) at 95% confidence. Further analysis with a simple logistic regression test showed that female adolescents who were past smokers were 12.59 times more likely (95% CI: 12.37–12.82) to experience depression than adolescents who were never smokers. Female adolescents who are current smokers have an 8.10 times greater chance (95% CI: 7.79–8.43) than female adolescents who were never smokers (Table 2).

The results of multivariate modeling showed that female adolescents who were ex-smokers were 10.93 times more likely (95% CI: 10.73–11.13) to experience depression compared to female adolescents who were never smokers. Meanwhile, female adolescents who are current smokers are 5.82 times more likely (95% CI 5.58–6.08) to experience depression compared to those who were never smokers. Further results were also obtained that divorced female adolescents were 4.32 times more likely (95% CI: 4.15–4.49) to experience depression than unmarried women. Female adolescents living in urban areas were 2.12 times more likely (95% CI: 2.10–2.13) to experience depression than those living in rural areas (Table 3).

## DISCUSSION

This study identified that the prevalence of depression is 1.6% among older female adolescents in Indonesia. This prevalence is lower than that of other national surveys in Indonesia. One study reported a prevalence of 6.7% for female adolescents aged 10–17 years<sup>6</sup>. Research in Indonesia using the Indonesia Family Life Survey data on female adolescents aged 15–19 years also showed much higher results, namely 26.3%, compared to those who were not depressed<sup>9</sup>. The differences in findings across prevalence rates can be influenced by differences in study design, variation in study populations, and the definition or instrument used to assess depression.

Important findings in this study show that there is a relationship between smoking status, marital status, location of residence, and depression in female adolescents, which is consistent with previous research<sup>8,9</sup>. This finding is in line with Lawrence Green's theory that the incidence of depression in female adolescents can be influenced by the interaction of various determinants, including predisposing factors, enabling factors, and reinforcing factors.

The findings of our study reinforce previous evidence showing a link between smoking status and depression in female adolescents<sup>10,11</sup>. In line with that, a study in Indonesia on female adolescents aged 15–19 years also identified smoking status as one of the factors related to common mental disorders<sup>12</sup>. This relationship is thought to be related to nicotine exposure<sup>13,14</sup>. Recent studies have

shown that the association between nicotine exposure and depressive symptoms is more meaningful in women with increased depressive symptoms at certain nicotine exposure levels<sup>13</sup>. In addition, it is thought that active maturation of the acetylcholine and dopamine systems that play a role in the regulation of mood, affect, and cognitive control in adolescents may be disrupted in their development by nicotine exposure associated with susceptibility to mental health disorders, including depression<sup>14,15</sup>. In line with this, previous studies have found that a younger age of first exposure to cigarettes is associated with an increased risk of major depressive episodes<sup>16,17</sup>.

In our analysis, ex-smokers were at a higher risk for depression than those who were never smokers, and the risk was even higher than that of current smokers. Research in Indonesia on adolescents and young adults is also in line with these results<sup>8,9</sup>. Consistent with this, a national study in the US found that ex-smokers were also associated with a higher risk of depression than non-smokers, but the amount of risk was no greater than that of current smokers<sup>18</sup>. In line with the study, the association between ex-smokers and depressive symptoms is likely to be higher in females; other factors, such as having recently quit smoking, having a history of high cumulative exposure, and former dual users, are also suspected to be related<sup>19</sup>. In women, the high risk of depression in ex-smokers is thought to be related to nicotine withdrawal symptoms that are influenced by fluctuations in ovarian hormones throughout the menstrual cycle<sup>20</sup>. However, given the cross-sectional design of this study, the direction of this relationship is uncertain, and there is a possibility of reverse causality between smoking and depression.

Our study found that divorced/widowed female adolescents were 4.32 times more at risk of depression than those who had never married. These results are in line with a study in the Philippines on adolescents that reported that separation, including living divorced or widowed, was associated with a 2.02-fold increased risk of depressive symptoms compared to unmarried status<sup>21</sup>. Studies on divorced single-parent women show that there is self-stigma in divorced single-parent women, which is associated with an increased risk of mental health distress through decreased self-esteem<sup>22</sup>. These results suggest that the breakdown of marital bonds is likely accompanied by psychosocial burdens that increase vulnerability to depression.

Living in urban areas is associated with a higher risk of depression than living in rural areas in female adolescents in Indonesia. These findings are in line with previous research in Bangladesh that showed that female adolescents living in urban non-slum areas are more at risk of depression than those living in rural areas<sup>23</sup>. Similarly, a national study in South Africa reported that the prevalence of depression in adolescents living in urban areas was higher than that of rural adolescents, at 14.64% and 9.40%<sup>24</sup>. However, these findings are inconsistent with studies in Korea, which

show that the proportion of moderate-severe depression symptoms is higher in rural than urban groups and confirm that the determinants of depression differ according to the context of residence<sup>25</sup>. Symptoms of depression in female adolescents living in urban areas are more associated with assessments of personal health conditions and pressure due to academic achievement demands, while in rural groups, they are related to rule violations and academic problems<sup>25</sup>. Depression in female adolescents is likely contextual, so the influence of urbanity on depression is likely mediated by social, academic, and environmental pressures that differ between countries. Further research needs to use a longitudinal design in order to better elucidate the direction of the relationship between residence and depression in female adolescents.

### Strengths and limitations

This study has some strengths. This study used a large national sample of Indonesian late adolescent females. The main focus is on female adolescents aged 15–19 years, so the analysis is more specific to groups that are often overlooked in general studies of adolescents. Furthermore, the findings that smoking status is associated with depression in female adolescents in this study may strengthen the evidence of a link between smoking status and depression in these populations. There are limitations in this study that should be considered when interpreting the results. First, cross-sectional designs do not allow for causal or two-way relationships between variables. Second, this study used secondary data, so other variables that may affect depression in female adolescents could not be controlled in this study.

### Implications

These findings have important implications for efforts to promote mental health in female adolescents, especially groups in more vulnerable social conditions in Indonesia. The results of the study show the need to improve early detection of depression and the development of prevention and intervention programs that are sensitive to the context of female adolescents' development. Furthermore, longitudinal research and studies with national samples that specifically target the adolescent girl population are needed to explain the direction of stronger relationships.

### CONCLUSIONS

The prevalence of depression among female adolescents aged 15–19 years in Indonesia was 1.6%. Smoking status, marital status, and location of residence were associated with depression in this population. Female adolescents who were ex-smokers and current smokers had higher odds of depression than those who were never smokers, with the highest odds observed among ex-smokers. Divorced/widowed female adolescents also had higher odds of depression than those who had never married, and those living in urban areas had higher odds of depression than

those living in rural areas. These findings indicate that older female adolescents with these characteristics may represent important subgroups for depression screening and prevention efforts in Indonesia.

### REFERENCES

1. Sawyer SM, Azzopardi PS, Wickremarathne D, Patton GC. The age of adolescence. *Lancet Child Adolesc Health*. 2018;2(3):223-228. doi:[10.1016/S2352-4642\(18\)30022-1](https://doi.org/10.1016/S2352-4642(18)30022-1)
2. World Health Organization. Mental health of adolescents; 2025. Accessed April 21, 2026. <https://www.who.int/news-room/fact-sheets/detail/adolescent-mental-health>
3. World Health Organization. Adolescent Health; 2025. Accessed April 21, 2026. <https://www.who.int/health-topics/adolescent-health>
4. Clayborne ZM, Varin M, Colman I. Systematic review and meta-analysis: Adolescent depression and long-term psychosocial outcomes. *J Am Acad Child Adolesc Psychiatry*. 2019;58(1):72-79. doi:[10.1016/j.jaac.2018.07.896](https://doi.org/10.1016/j.jaac.2018.07.896)
5. Chang K, Kuhlman KR. Adolescent-onset depression is associated with altered social functioning into middle adulthood. *Sci Rep*. 2022;12(1):17320. doi:[10.1038/s41598-022-22131-1](https://doi.org/10.1038/s41598-022-22131-1)
6. Center for Reproductive Health, Universitas Gadjah Mada, Johns Hopkins Bloomberg School of Public Health, University of Queensland, Universitas Sumatera Utara, Universitas Hasanuddin. Indonesia National Adolescent Mental Health Survey (I-NAMHS) Report. Center for Reproductive Health, Universitas Gadjah Mada; 2023. Accessed April 21, 2026. <https://publichealth.jhu.edu/sites/default/files/2025-05/I-NAMHS-Report-English.pdf>
7. Health Development Policy Agency, Ministry of Health of the Republic of Indonesia. Indonesian Health Survey 2023. Ministry of Health of the Republic of Indonesia; 2023. Accessed April 21, 2026. <https://www.badankebijakan.kemkes.go.id/en/ski-2023-dalam-angka/>
8. Suryaputri IY, Mubasyiroh R, Idaiani S, Indrawati L. Determinants of depression in Indonesian Youth: Findings from a community-based survey. *J Prev Med Pub Health*. 2022;55(1):88-97. doi:[10.3961/jpmph.21.113](https://doi.org/10.3961/jpmph.21.113)
9. Purborini N, Lee MB, Devi HM, Chang HJ. Associated factors of depression among young adults in Indonesia: A population-based longitudinal study. *J Formos Med Assoc*. 2021;120(7):1434-1443. doi:[10.1016/j.jfma.2021.01.016](https://doi.org/10.1016/j.jfma.2021.01.016)
10. Yue Y, Hong L, Guo L, et al. Gender differences in the association between cigarette smoking, alcohol consumption, and depressive symptoms: A cross-sectional study among Chinese adolescents. *Sci Rep*. 2015;5(1):17959. doi:[10.1038/srep17959](https://doi.org/10.1038/srep17959)
11. Tomita A, Manuel JI. Evidence on the association between cigarette smoking and incident depression from the South African National Income Dynamics Study 2008–2015: Mental health implications for a resource-limited setting. *Nicotine Tob Res*. 2020;22(1):118-123. doi:[10.1093/ntr/nty163](https://doi.org/10.1093/ntr/nty163)
12. Asrullah M, L'Hoir M, Paulo MJ, Feskens EJM, Melse-Boonstra

- A. Determinants of Common Mental Disorders (CMD) among adolescent girls aged 15-19 years in Indonesia: Analysis of the 2018 National Basic Health Survey Data. *PLOS Glob Public Health*. 2022;2(3):e0000232. doi:[10.1371/journal.pgph.0000232](https://doi.org/10.1371/journal.pgph.0000232)
13. Liu Y, Zhang L, Fu S, Wei S, Jin Z, He L. Gender differences in the relationship between nicotine exposure and symptoms of depression. *Pharmacol Biochem Behav*. 2024;244:173857. doi:[10.1016/j.pbb.2024.173857](https://doi.org/10.1016/j.pbb.2024.173857)
  14. Reynolds LM, Faure P, Barik J. Adolescent nicotine exposure and persistent neurocircuitry changes: Unveiling lifelong psychiatric risks. *Mol Psychiatry*. 2025;30(11):5534-5545. doi:[10.1038/s41380-025-03110-0](https://doi.org/10.1038/s41380-025-03110-0)
  15. Ng THJ, Sarikahya MH, Hudson R, et al. Adolescent nicotine exposure induces long-term, sex-specific disturbances in mood and anxiety-related behavioral, neuronal, and molecular phenotypes in the mesocorticolimbic system. *Neuropsychopharmacology*. 2024;49(7):1171-1182. doi:[10.1038/s41386-024-01853-y](https://doi.org/10.1038/s41386-024-01853-y)
  16. Ranjit A, Buchwald J, Latvala A, et al. Predictive association of smoking with depressive symptoms: A longitudinal study of adolescent twins. *Prev Sci*. 2019;20(7):1021-1030. doi:[10.1007/s11121-019-01020-6](https://doi.org/10.1007/s11121-019-01020-6)
  17. Devkota B, Salas J, Garfield L. Increased risk of major depression with early age of exposure to cigarettes. *Am J Prev Med*. 2016;51(6):933-938. doi:[10.1016/j.amepre.2016.05.022](https://doi.org/10.1016/j.amepre.2016.05.022)
  18. Wu Z, Yue Q, Zhao Z, et al. A cross-sectional study of smoking and depression among US adults: NHANES (2005–2018). *Front Public Health*. 2023;11:1081706. doi:[10.3389/fpubh.2023.1081706](https://doi.org/10.3389/fpubh.2023.1081706)
  19. Kim J, Lee E, Hwang D, Ko AJ, Park EC. Association between smoking cessation and depressive symptoms according to cessation duration, pack-years, and tobacco product type: A nationwide cross-sectional study in Korea. *Front Public Health*. 2026;14:1755259. doi:[10.3389/fpubh.2026.1755259](https://doi.org/10.3389/fpubh.2026.1755259)
  20. Giner P, Ortegon S, Bagdas D, O'Dell LE. The influence of ovarian hormones on the putative mechanisms that promote female nicotine use. *Curr Opin Neurobiol*. 2024;88:102900. doi:[10.1016/j.conb.2024.102900](https://doi.org/10.1016/j.conb.2024.102900)
  21. Puyat JH, Salvador DL, Tuazon AC, Afable SD. Rising prevalence of depression and widening sociodemographic disparities in depressive symptoms among Filipino youth: Findings from two large nationwide cross-sectional surveys. *Camb Prisms Glob Ment Health*. 2025;12:e51. doi:[10.1017/gmh.2025.39](https://doi.org/10.1017/gmh.2025.39)
  22. Kim A, Jeon S, Song J. Self-stigma and mental health in divorced single-parent women: Mediating effect of self-esteem. *Behav Sci*. 2023;13(9):744. doi:[10.3390/bs13090744](https://doi.org/10.3390/bs13090744)
  23. Mridha MK, Hossain MM, Khan MSA, et al. Prevalence and associated factors of depression among adolescent boys and girls in Bangladesh: Findings from a nationwide survey. *BMJ Open*. 2021;11(1):e038954. doi:[10.1136/bmjopen-2020-038954](https://doi.org/10.1136/bmjopen-2020-038954)
  24. Ajaero CK, Nzeadibe CT, Igboeli EE. Rural-urban differences in the prevalence and predictors of depression among adolescents in South Africa. *South Afr J Child Health*. 2018;12(2b):71. doi:[10.7196/SAJCH.2018.v12i2b.1509](https://doi.org/10.7196/SAJCH.2018.v12i2b.1509)
  25. Lee G, Ham OK, Lee BG, Kim AM. Differences in factors associated with depressive symptoms between urban and rural female adolescents in Korea. *J Korean Acad Nurs*. 2018;48(4):475. doi:[10.4040/jkan.2018.48.4.475](https://doi.org/10.4040/jkan.2018.48.4.475)

#### ACKNOWLEDGMENTS

The authors would like to express their gratitude to the Health Development Policy Agency (BKPK), Ministry of Health of the Republic of Indonesia, for permission to use secondary data from the 2023 Indonesian Health Survey (SKI). Appreciation is also extended to the entire survey team, enumerators, and respondents who contributed to the collection of these nationwide health data. Access to these data has been instrumental in efforts to develop mental health policies for female adolescents in Indonesia. The authors also gratefully acknowledge the support and encouragement provided by the Faculty of Medicine and Health Sciences (FKIK), Universitas Bengkulu.

#### CONFLICTS OF INTEREST

The authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none was reported.

#### FUNDING

There was no source of funding for this research.

#### ETHICAL APPROVAL AND INFORMED CONSENT

Ethical approval and informed consent were not required for this study as it is a secondary analysis of existing data.

#### DATA AVAILABILITY

The data supporting this research are available from the authors on reasonable request.

#### PROVENANCE AND PEER REVIEW

Not commissioned; externally peer-reviewed.